

Vaccines for the National Immunisation Schedule



A reference card for vaccinators and other health professionals

our best protection

<p>Pregnancy</p>	 <p>Tdap injection (Boostrix[®])</p>	<p>+</p>	<p>Influenza 1 injection (given annually) Brand varies.</p>	
<p>6 Weeks</p>	 <p>RV1 oral vaccine (Rotarix[®])</p>	<p>+</p>	 <p>DTaP-IPV-Hep B/Hib injection (Infanrix[®] hexa)</p>	<p>+</p>  <p>PCV13 injection (Prevenar 13[®])</p>
<p>3 Months</p>	 <p>RV1 oral vaccine (Rotarix[®])</p>	<p>+</p>	 <p>DTaP-IPV-Hep B/Hib injection (Infanrix[®] hexa)</p>	<p>+</p>  <p>+ MenB injection (Bexsero[®]) Can be given at 8 weeks</p>
<p>5 Months</p>	 <p>DTaP-IPV-Hep B/Hib injection (Infanrix[®] hexa)</p>	<p>+</p>	 <p>PCV13 injection (Prevenar 13[®])</p>	<p>+</p>  <p>+ MenB injection (Bexsero[®]) Can be given at 4 months</p>
<p>12 Months</p>	 <p>MMR injection (Priorix[®])</p>	<p>+</p>	 <p>PCV13 injection (Prevenar 13[®])</p>	<p>+</p>  <p>+ MenB injection (Bexsero[®])</p>
<p>15 Months</p>	 <p>Hib injection (Hiberix[®])</p>	<p>+</p>	 <p>MMR injection (Priorix[®])</p>	<p>+</p>  <p>Varicella injection (Varivax[®])</p>
<p>4 Years</p>	 <p>DTaP-IPV injection (Infanrix[®] IPV)</p>			
<p>11+12 Years</p>	 <p>Tdap injection (Boostrix[®])</p>	<p>+</p>	 <p>HPV injection (Gardasil[®]9) (2 doses, 6 months apart)</p>	
<p>45 Years</p>	 <p>Tdap injection (Boostrix[®])</p>			
<p>65 Years</p>	 <p>Tdap injection (Boostrix[®])</p>	<p>+</p> <p>Influenza 1 injection (given annually). Brand varies.</p>	<p>+</p>  <p>Shingles injection (Shingrix[®])</p>	

The National Immunisation Schedule

Age	Disease to protect against	Vaccine
Pregnancy	Tetanus + diphtheria + whooping cough (pertussis)	Boostrix [®]
	Influenza	Brand varies.
6 Weeks	Rotavirus (first dose must be given before 15 weeks)	Rotarix [®] (oral)
	Diphtheria + tetanus + whooping cough (pertussis) + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	Infanrix [®] hexa
	Pneumococcal disease	Prevenar 13 [®]
3 Months	Rotavirus (second dose must be given before 25 weeks)	Rotarix [®] (oral)
	Diphtheria + tetanus + whooping cough + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	Infanrix [®] hexa
	Meningococcal B (can be given at 8 weeks)	Bexsero [®]
5 Months	Diphtheria + tetanus + whooping cough + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	Infanrix [®] hexa
	Pneumococcal disease	Prevenar 13 [®]
	Meningococcal B (can be given at 4 months)	Bexsero [®]
12 Months	Measles + mumps + rubella	Priorix [®]
	Pneumococcal disease	Prevenar 13 [®]
	Meningococcal B	Bexsero [®]
15 Months	<i>Haemophilus influenzae</i> type b (Hib)	Hiberix [®]
	Measles + mumps + rubella	Priorix [®]
	Chickenpox (varicella)	Varivax [®]
4 Years	Diphtheria + tetanus + whooping cough + polio	Infanrix [®] IPV
11+12 Years	Tetanus + diphtheria + whooping cough	Boostrix [®]
	Human papillomavirus (HPV)	Gardasil [®] 9 (2 doses, 6 months apart)
45 Years	Tetanus + diphtheria + whooping cough	Boostrix [®]
65 Years	Tetanus + diphtheria + whooping cough	Boostrix [®]
	Influenza	Brand varies.
	Shingles	Shingrix [®]